

Appendix C: BW Agent Characteristics

Disease	Transmit Man to Man	Infective Dose (Aerosol)	Incubation Period	Duration of Illness	Lethality (approx. case fatality rates)	Persistence of Organism	Vaccine Efficacy (aerosol exposure)
Inhalation anthrax	No	8,000-50,000 spores	1-6 days	3-5 days (usually fatal if untreated)	High	Very stable - spores remain viable for > 40 years in soil	2 dose efficacy against up to 1,000 LD ₅₀ in monkeys
Brucellosis	No	10 -100 organisms	5-60 days (usually 1-2 months)	Weeks to months	<5% untreated	Very stable	No vaccine
Cholera	Rare	10-500 organisms	4 hours - 5 days (usually 2-3 days)	≥ 1 week	Low with treatment, high without	Unstable in aerosols & fresh water; stable in salt water	No data on aerosol
Glanders	Low	Assumed low	10-14 days via aerosol	Death in 7-10 days in septicemic form	> 50%	Very stable	No vaccine
Pneumonic Plague	High	100-500 organisms	2-3 days	1-6 days (usually fatal)	High unless treated within 12-24 hours	For up to 1 year in soil; 270 days in live tissue	3 doses not protective against 118 LD ₅₀ in monkeys
Tularemia	No	10-50 organisms	2-10 days (average 3-5)	≥ 2 weeks	Moderate if untreated	For months in moist soil or other media	80% protection against 1-10 LD ₅₀
Q Fever	Rare	1-10 organisms	10-40 days	2-14 days	Very low	For months on wood and sand	94% protection against 3,500 LD ₅₀ in guinea pigs
Smallpox	High	Assumed low (10-100 organisms)	7-17 days (average 12)	4 weeks	High to moderate	Very stable	Vaccine protects against large doses in primates
Venezuelan Equine Encephalitis	Low	10-100 organisms	2-6 days	Days to weeks	Low	Relatively unstable	TC 83 protects against 30-500 LD ₅₀ in hamsters
Viral Hemorrhagic Fevers	Moderate	1-10 organisms	4-21 days	Death between 7-16 days	High for Zaire strain, moderate with Sudan	Relatively unstable - depends on agent	No vaccine
Botulism	No	0.001 µg/kg is LD ₅₀ for type A	1-5 days	Death in 24-72 hours; lasts months if not lethal	High without respiratory support	For weeks in nonmoving water and food	3 dose efficacy 100% against 25-250 LD ₅₀ in primates
Staph Enterotoxin B	No	0.03 µg/person incapacitation	3-12 hours after inhalation	Hours	< 1%	Resistant to freezing	No vaccine
Ricin	No	3-5 µg/kg is LD ₅₀ in mice	18-24 hours	Days - death within 10-12 days for ingestion	High	Stable	No vaccine

T-2 Mycotoxins	No	Moderate	2-4 hours	Days to months	Moderate	For years at room temperature	No vaccine
---------------------------	----	----------	-----------	----------------	----------	-------------------------------	------------